

Ferrocenylalkyl carbocations. II. Contribution to the problem of the structure of ferrocenylmethyl carbocation

Ursić, S.

Source / Izvornik: **Croatica Chemica Acta, 1982, 55, 305 - 308**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:163:632526>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-17**



Repository / Repozitorij:

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



CCA-1334

YU ISSN 0011-1643

UDC 546.73:541.124

Preliminary communication

Ferrocenylalkyl Carbocations. II. Contribution to the Problem of the Structure of Ferrocenylmethyl Carbocation

S. Uršić

Faculty of Pharmacy and Biochemistry, University of Zagreb, 41000 Zagreb, Croatia, Yugoslavia

Received November 2, 1981

In this communication, an unusually large secondary α -deuterium kinetic isotope effect for hydrolysis of the ferrocenylmethylquinolinium ion, that elucidates significantly the problem of the structure and manner in which stabilisation of the α -ferrocenylalkyl carbocations occurs, will be reported. This problem arises mainly from uncertainties as to whether some direct bonding between iron and the cationic center is of significance in the stabilisation of α -ferrocenylalkyl carbocations¹.

Hydrolysis of ferrocenylmethylquinolinium chloride was carried out at 25 °C. Rate constants for hydrolysis in water and 80:20 water-acetonitrile were determined spectrophotometrically. Results, together with corresponding α -deuterium kinetic isotope effects are given in Table I.

Some secondary α -deuterium kinetic isotope effects for C—N bond cleavage in unimolecular solvolysis have been determined²⁻⁴, (Table II) and only a few of them are similar in magnitude to the isotope effect reported here. The magnitude of a kinetic isotope effect is directly connected with the transition state structure, and variations in the transition state structure should result in variations in the isotope effect. Cordes et al. suggested² that very similar α -deuterium isotope effects obtained in inosine and adenosine hydrolysis, (first and second entry in Table II) originate from a C—N bond being largely or completely cleaved in the transition state. Similarly, an α -deuterium isotope effect of ca. 1.24 per deuterium atom for hydrolysis of the ferrocenylmethylquinolinium ion in water clearly suggested that the $sp^3 \rightarrow sp^2$ rehybridization is nearly complete or complete on going from reactants to the transition state. Such a conclusion is supported if one considers the similarity in secondary α -deuterium kinetic isotope effects for C—O and C—N bond cleavage^{1,4}. Thus, an α -deuterium isotope effect of ca. 1.23 is usually considered to be near to the maximum expected for C—O bond cleavage⁵.

The present results for hydrolysis of the ferrocenylquinolinium ion in water strongly indicate that there is no significant bonding between iron and the α -cationic center in the transition state for hydrolysis, since such a participation of iron must result in an effect substantially smaller than the isotope effect observed^{4,5}. Therefore, most of the models^{6a} for α -ferrocenylalkyl carbocation which allow direct bonding between iron and α -carbon probably need to be

TABLE I
Rates of Hydrolysis of Ferrocenylmethylquinolinium Ion at 25 °C

Substrate ^a	Solvent	Rate constant ^{a,e} 10 ⁴ s ⁻¹	α -deuterium kinetic isotope effect
FcCH ₂ Q ⁺	Water ^b	0.658 ± 0.006	1.479 ± 0.024
FcCD ₂ Q ⁺		0.445 ± 0.006	
FcCH ₂ Q ⁺	Water- acetonitrile ^c 80 : 20 v/v	0.798 ± 0.020	1.237 ± 0.049
FcCD ₂ Q ⁺		0.645 ± 0.020	

^a) Fc = ferrocenyl, Q = quinolinium. Ferrocenylmethylquinolinium chloride was prepared from ferrocenylmethyl carbinol and quinolinium hydrochloride in acetonitrile. Corresponding deuterated substance was prepared analogously from deuterated carbinol (see ref. 1). M. p. 119–121 °C. Assay of FcCH₂QCl 99% (by potentiometric titration of chloride). Log ϵ (in water) is 4.44(236 nm), 3.45(280 nm), 3.76(313 nm), 2.13(380 nm) and 2.47(443 nm). Spectra of solvolytic products in comparison with spectra of authentic samples (ferrocenylcarbinol and quinolinium hydrochloride) confirmed the structural assignment. When ferrocenylcarbinol in solvolysis mixture was (after 8 half-lives) oxidized with ferric perchlorate to the corresponding ferricenium product, comparison of absorbance at 627 nm with an authentic sample of oxidized carbinol indicated (within experimental error) that solvolysis of FcCH₂Q⁺ gives quantitatively ferrocenylcarbinol.

^b) With 0.3% v/v acetonitrile added. Ionic strength of 0.5 was maintained with benzyltrimethylammonium chloride.

^c) Ionic strength 0.3, salt benzyltrimethylammonium chloride.

^d) Kinetics were measured spectrophotometrically, by following differences in absorbance at 456 nm. Individual rate constants were computed using a nonlinear regression analysis. Reported rate constants are the mean of four (hydrolysis in water) or eight (hydrolysis in water-acetonitrile) individual constants.

^e) Control experiments with added base (NaOH, amount equal to that of FcCH₂Q⁺ substrate) or acid (10⁻³ M dm⁻³ HCl, fivefold acid over the substrate concentration) shows no significant rate depression or rate acceleration. In addition, almost identical rate constants were obtained from data for one half-life and assumed infinity absorbance (from spectra of quinoline hydrochloride and ferrocenylmethyl carbinol) and from data for four half-lives of solvolysis.

TABLE II
Kinetic Secondary α -Deuterium Isotope Effects for C—N Bond Cleavage

Substrate	Leaving group	Kinetic α -deuterium Isotope effect	Ref.
inosine	hypoxantyne ^a	1.23 ± 0.02	2
adenosine	adenine ^a	1.229 ± 0.012	2
β -D-galactopyranosyl-4-bromoisquinolinium ion	4-bromoisquinoline	1.187 ± 0.046	3a
β -D-galactopyranosylisoquinolinium ion	isoquinoline	1.17 ± 0.03	3b
β -D-galactopyranosylpyridinium ion	pyridine	1.13 ± 0.04 ^b	3c
ferrocenylmethyl-trimethylammonium ion	trimethylamine	1.06 ± 0.04 ^{c,d}	4
ferrocenylmethylquinolinium ion	quinoline	1.479 ± 0.024 ^c	this work

^a) purine base or protonated purine base

^b) At 100 °C, corresponding to one isotope effect of 1.16 at 25 °C

^c) Isotope effects from two α -deuterium atoms

^d) at 80 °C

retracted. The ferrocenylmethyl carbocation was isolated as its tetrafluoroborate⁷, but X-ray crystal analysis was performed only for some related secondary and tertiary carbocations⁸. These analyses offer some possibility for iron-cationic center interaction. Perhaps, there exist some differences between the transition state structure for solvolysis and the structure of α -ferrocenylalkyl carbocations in solid state and in solution.*

In 80 : 20 water-acetonitrile there is an essentially smaller α -deuterium isotope effect for solvolysis and somewhat greater rate of hydrolysis of an ferrocenylmethylquinolinium ion (cf. third and fourth entry in Table I). The small rate enhancement for hydrolysis is not surprising. This enhancement is expected for solvolysis of the cationic substrate when the dielectric constant of solvent is decreased, due to dispersal of charge in the transition state⁹. The magnitude of an α -deuterium isotope effect of ca. 1.12 (Table I) per deuterium atom is similar to that for hydrolysis of the β -D-galactopyranosylpyridinium ion (fifth entry in Table II), and to the α -deuterium kinetic isotope effect k_D/k_H calculated earlier for addition of pyridine to the ferrocenylmethyl carbocation in water¹.

It was observed¹⁰ that addition of organic solvent to water markedly destabilised ferrocenylalkyl carbocations. On the other hand, Sinott and Jencks¹¹ reported that there is some specific but weak interaction with both the leaving group and the incoming solvent molecule in the transition state for solvolysis of α -D-glucopyranosylpyridinium ions. These interactions may be predominantly or entirely electrostatic.

There is probably, some interaction with solvent molecules in the transition state in the case of hydrolysis of ferrocenylmethylquinolinium ion in water-acetonitrile, which causes diminishing of the α -deuterium isotope effect. This problem will be further elaborated in an subsequent paper.

Acknowledgement: — The author wishes to thank dr. M. Biruš, Faculty of Pharmacy and Biochemistry, for allowing the use of his computer programs and M. Essert, Faculty of Engineering and Shipbuilding at the University of Zagreb, for help and computer time on the PDP-11 computer sistem. Acknowledgement is made to the Croatian Research Council for financial support.

REFERENCES

1. S. Uršić, *Croat. Chem. Acta* **54** (1981) 69 and references cited therein.
2. R. Romero, R. Stein, H. G. Bull, and E. H. Cordes, *J. Amer. Chem. Soc.* **100** (1978) 7620.
3. a) M. L. Sinott and S. G. Withers, *Biochem. J.* **143** (1974) 571; M. L. Sinott, O. M. Viratelle, and S. G. Withers, *Biochem. Soc. Trans.* **3** (1975) 1005; c) C. C. Jones, M. L. Sinott, and I. J. L. Souchar, *J. Chem. Soc. Perkin II* (1977) 1191.
4. S. Uršić and S. Ašperger, *J. Inorg. Nucl. Chem.* **41** (1979) 1329.
5. V. J. Shiner, Jr. in *Isotope Effects in Chemical Reactions* eds. C. J. Collins and N. S. Bowman, van Nostrand Reinhold, New York 1970, Ch. 2.
6. a) For discussion concerning models of α -ferrocenylalkyl ions see for example, A. A. Koridze, P. V. Petrovskii, S. P. Gubin, V. I. Sokolov, and A. I. Mokhov, *J. Organometal. Chem.* **136** (1977) 65.
b) G. Neshvad, R. M. G. Roberts, and J. Silver, *J. Organometal. Chem.* **221** (1981) 85.
7. A. Cecon, G. Giacometti, A. Venzo, D. Paolucchi, and D. Benozzi, *J. Organometal. Chem.* **185** (1980) 231.

* Recent Mössbauer spectroscopy analysis^{6b} of α -ferrocenylalkyl carbocations seems to indicate some direct iron-cationic center interaction for FcC^+H_2 in frozen acidic solution.

8. M. Cais, S. Dani, F. H. Herbststein, and M. Kapon, *J. Amer. Chem. Soc.* **100** (1978) 5554; U. Behrens, *J. Organometal. Chem.* **182** (1979) 89.
9. Similar results were obtained for solvolysis of 2-ferrocenylethyltrimethylammonium iodide, and some sulfonium salts. See: G. Gokel, D. Marquarding, and I. Ugi, *J. Org. Chem.* **37** (1972) 3052 and references cited therein.
10. C. A. Bunton, N. Carrasco, and W. E. Watts, *J. Chem. Soc. Perkin II* (1979) 1267.
11. M. L. Sinott and W. P. Jencks, *J. Amer. Chem. Soc.* **102** (1980) 2026.

SAŽETAK

Ferocenilalkil karbokationi. II. Doprinos problemu strukture ferocenilmetil karbokationa

Stanko Uršić

Odreden je sekundarni α -deuterijski kinetički izotopni efekt u solvolizi ferocenilmetilkinolinijeva iona i iznosi ca. 24% po atomu deuterija kod solvolize u vodi, te ca. 12% po D atomu kod solvolize u 80/20 vodi/acetonitrilu. Diskutirano je značenje tih rezultata u vezi sa strukturom ferocenilmetil karbokationa.