

# Reactions with 1-benzotriazolecarboxylic acid chloride. V Synthesis of sulfonylureas and related compounds

---

Butula, I .; Vela, V .; Proštenik, M. V.

Source / Izvornik: **Croatica Chemica Acta, 1979, 52, 47 - 49**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:163:149607>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-09-13**



Repository / Repozitorij:

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



CCA-1142

YU ISSN 0011-1643

547.495.2:496

Note

## Reactions with 1-Benzotriazolecarboxylic Acid Chloride. V<sup>1</sup>. Synthesis of Sulfonylureas and Related Compounds

I. Butula, \*V. Vela, and \*M. V. Proštenik

Faculty of Pharmacy and Biochemistry, University of Zagreb and \*PLIVA,  
Pharmaceutical and Chemical Works, 41001 Zagreb, Croatia, Yugoslavia

Received November 27, 1978


Sulfonylureas and related compounds, with hypoglycemic activity were synthesized, starting from corresponding sulfonylamide salts and 1-carbamoylbenzotriazoles.

Usually, sulfonylureas I with hypoglycemic activity were prepared in two commonly used ways. The first is based on the reaction of arylsulfonyl isocyanates<sup>2</sup> or arylsulfonyl carbamates<sup>3-6</sup> with amines, and the second on the reaction of sulfonylamide salts with isocyanates<sup>6-8</sup> or amidocarbonic acid derivatives<sup>2,6</sup>.

As a part of our work on the application of 1-carbamoylbenzotriazole derivatives II in organic synthesis<sup>9</sup>, we have recently described it as a very useful reagent for preparing ureas<sup>1</sup> in the reaction with primary and secondary amines. The reaction proceeded by means of direct nucleophilic attack of the amino group on the carbonyl, activated by a benzotriazole substituent.






In order to test the generality of the observed behaviour of II in the reaction with amines, the reaction with sulfonylamides III was investigated. The reaction was found to be a convenient route to sulfonylureas I, with regard to the useful properties of carbamoylbenzotriazoles, stable solids, which could be handled more easily with safety than isocyanate or phosgene, and because of the high yields obtained.

Temperatures (about 100 °C) for the reaction of sulfonylamides III, at which there was no evidence of dissociation of II into benzotriazole and isocyanate (as checked by IR spectroscopy of II in dioxane solution at 80 °C) indicated a direct nucleophilic attack of the sulfonylamide amino group on carbonyl, following the mechanism proposed for the formation of ureas<sup>1</sup>.



The reaction was carried out in dioxane or dimethylformamide solution at elevated temperature for 1 to 5 h. The reaction mixture was evaporated to dryness at reduced pressure. Sulfonylureas I were isolated by acidification of the water solution of the residue.

TABLE  
Preparation of Sulfonylureas Ia—Ie

Compound	R <sup>1</sup>	R <sup>2</sup>	Reaction conditions solvent/time/h	temp./°C	Method of isolation	Yield %	m. p./°C	lit. m. p./°C
Ia	CH <sub>3</sub>	CH <sub>3</sub>	DMF / 1	95	A	67	168—170	170—172 <sup>10</sup>
Ib	CH <sub>3</sub>	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	Dioxane / 1	100	A	79	148—150	141—143 <sup>10</sup>
Ic	CH <sub>3</sub>	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	DMF / 2	95	A	62	122—126	127—129 <sup>10</sup>
Id	CH <sub>3</sub>	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	Dioxane / 1	100	A	93	167—169	170—173 <sup>10</sup>
Ie	Cl	CH <sub>3</sub>	Dioxane / 1	100	A	62	186—190	188—189 <sup>10</sup>
If	Cl	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	Dioxane / 4	100	A	65	128—130	126—128 <sup>10</sup>
Ig	Cl	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	Dioxane / 1	100	A	89	115—117	115—116 <sup>10</sup>
Ih	Cl	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	Dioxane / 1	100	B	62	155—157	158—159 <sup>10</sup>
Ii		<i>c</i> -C <sub>6</sub> H <sub>11</sub>	Dioxane / 5	100	B	76	168—172	168—172 <sup>8</sup>
Ij			Dioxane / 4	100	B	80	190	189 <sup>6b</sup>
Ik		<i>c</i> -C <sub>6</sub> H <sub>11</sub>	Dioxane / 1	100	A	75	186—188	184 <sup>5</sup>
II	CH <sub>3</sub>		Dioxane / 5	100	B	65	168—171	171—173 <sup>2</sup>

## EXPERIMENTAL

*Preparation of Sulfonylureas I: General Procedure*

Sulfonylamide sodium salt III (5 mmol), 1-carbamoylbenzotriazole II (5 mmol) and an appropriate solvent (30 ml) were heated at 95–100 °C and the sulfonylurea thus formed was isolated by two methods:

*Method A.* — The solvent was evaporated in vacuo and 15% hydrochloric acid was added. After standing in a water bath at 60 °C for 0.5 h the product was filtered off.

*Method B.* — The insoluble part of the reaction mixture was filtered off, dissolved in 75 ml of water and the product was precipitated by the addition of 5% hydrochloric acid (to pH 3–4).

## REFERENCES

1. I. Butula, V. Vela, and B. Ivezić, *Croat. Chem. Acta* **51** (1978) 339.
2. J. B. Wright, *German Patent* 1,186,865; *C. A.* **63** (1965) 14760.
3. F. J. Marshall and M. V. Sigal, *J. Org. Chem.* **23** (1958) 927.
4. K. Grezon, E. V. Krumkalns, R. L. Brindle, F. J. Marshall, and M. A. Root, *J. Med. Chem.* **6** (1963) 760.
5. H. Weber, W. Aumüller, R. Weyer, K. Muth and F. H. Schmidt, *German Patent* 1,185,180; *C. A.* **62** (1965) 13092.
6. a) J. B. Wright and R. F. Willette, *J. Med. Pharm. Chem.* **5** (1962) 815.  
b) H. Plumpe and W. Puls, *US Patent* 3,755,587; *C. A.* **79** (1973) 115564.
7. *Holland Patent* 6,610,580; *C. A.* **68** (1968) 12733.
8. H. Weber, W. Aumüller, R. Weyer, and H. Schmidt, *German Patent* 1,283,837; *C. A.* **70** (1969) 47140.
9. I. Butula, M. V. Proštenik, and V. Vela, *Croat. Chem. Acta* **49** (1977) 837.
10. H. Rushing, G. Korger, W. Aumüller, H. Wagner, R. Weyer, A. Bänder, and J. Scholz, *Arzeim. Forsch.* **8** (1958) 448.

## SAŽETAK

**Reakcije klorida 1-benzotriazolkarboksilne kiseline. V. Sinteza sulfonylurea i srodnih spojeva**

*I. Butula, \*V. Vela i \*M. V. Proštenik*

Opisana je priprava sulfonylurea i srodnih spojeva koji imaju hipoglikemijsko djelovanje. U sintezi se polazi od soli sulfonylamida i 1-karbamoilbenzotriazola.

FARMACEUTSKO-BIOKEMIJSKI FAKULTET

i

Prispjelo 27. studenog 1978.

\*SOUR-PLIVA, FARMACEUTSKA,  
KEMIJSKA, PREHRAMBENA I KOZMETIČKA  
INDUSTRIJA, 41000 ZAGREB