

Kinetic study of the substitution of pyridine by cyanide in the bis(pyridine)cobalt(III)hematoporphyrin-IX: Distinguishing between the Id and D mechanism

Biruš, M.; Pribanić, M.

Source / Izvornik: *Croatica Chemica Acta*, 1977, 49, 65 - 68

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:163:389119>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom](#).

Download date / Datum preuzimanja: **2024-07-19**



Repository / Repozitorij:

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



Kinetic Study of the Substitution of Pyridine by Cyanide in the bis(pyridine)cobalt(III)hematoporphyrin-IX: Distinguishing between the I_d and D Mechanism


M. Biruš and M. Pribanić*

Department of Chemistry, Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia, Yugoslavia

Received August 11, 1976

A kinetic study of the reaction between the cyanide ion and the bis(pyridine)cobalt(III)hematoporphyrin-IX complex to give (CN)₂cobalt(III)hematoporphyrin-IX has shown that the axial ligand substitution occurs by a purely dissociative (D but not I_d) mechanism in chloroform involving predissociation to a five-coordinate (pyridine)cobalt(III)hematoporphyrin-IX complex.

A dissociative mechanism has been advanced for most of the metalloporphyrin substitution reactions.^{1,2} It is surprising that »mass-law (rate) retardation« in noncoordinating solvents has been utilized only to a minor extent for differentiation between closely related I_d and D mechanisms.^{3,4} Our kinetic studies of the reaction of the cyanide ion with (py)₂Co(hemato-IX) (py = pyridine, hemato-IX = hematoporphyrin-IX) in chloroform eliminate the possibility of the I_d process and provide strong support for the D mechanism.** The stoichiometry of the over-all reaction was found spectrophotometrically to be:




where the reactant and product were identified by their visible absorption spectrum⁵.

Reaction (1) proceeds to completion under the conditions employed in this study. The kinetics were studied spectrophotometrically by monitoring the change in the Soret region at 25 °C in chloroform containing less than 0.01 M [H₂O] and under pseudo-first-order conditions. Our kinetic data show an inverse dependence on pyridine and between the first- and zero-order in cyanide ion concentration. This observation supports the previous statement that the second axial anion incorporation is rapid.^{1,2,6} The data presented are in full agreement with the dissociative mechanism depicted by equation (2).

* The correspondence should be addressed to this author.

** The over-all charge on the metalloporphyrin molecule is not shown throughout the paper.



The rate law, assuming a steady state for the five-co-ordinate intermediate (py)Co(hemato-IX) is

$$\text{rate} = \frac{k_1 [(\text{py})_2\text{Co(hemato-IX)}]}{(k_{-1}/k_2) ([\text{py}]/[\text{CN}^-]) + 1} = k_{\text{obs}} [(\text{py})_2\text{Co(hemato-IX)}] \quad (3)$$




Figure $(k_{\text{obs}})^{-1}$ vs. concentration ratio [pyridine]/[CN⁻] for reaction (1).

The plot of $(k_{\text{obs}})^{-1}$ versus $[\text{py}]/[\text{CN}^-]$ gives a straight line with a nonzero intercept and yield the k_{-1}/k_2 and k_1 values of 0.35 and $2.50 \times 10^{-3} \text{ s}^{-1}$, respectively.

An obvious variant of this mechanism, which is kinetically indistinguishable from that shown above, is the one which involves formation of the intermediate aquo-complex *via* the I_d mechanism, since the solvent chloroform may contain as much as 0.01 M [H₂O]. The application of the steady state condition to the intermediate aquo-complex leads to a rate law identical in form with equation (3), but it can be ruled out on the basis of the fact that the controlled varying water amount in chloroform does not influence the rate of axial substitution as shown in the Table. This conclusion is unsafe if

the cation-dipol association constant between the substrate and water molecule is very high.

The extension of this method to other bis-ligated metalloporphyrins is in progress.

TABLE
Rate Data for the Reaction (2)

[py]/mol l ⁻¹	[CN ⁻] ^a /mol l ⁻¹	k _{obs} /s ⁻¹	k ₋₁ /k ₂
0	0.08	2.7 × 10 ⁻³	
0	0.02	2.1 × 10 ⁻³	
0	0.005	2.4 × 10 ⁻³	
0	0.004	2.5 × 10 ⁻³	
0	0.004	2.4 × 10 ^{-3b}	
0	4 × 10 ⁻⁴	2.7 × 10 ⁻³	
4 × 10 ⁻³	2.5 × 10 ⁻⁴	3.9 × 10 ⁻⁴	0.34
8 × 10 ⁻⁴	2.5 × 10 ⁻⁴	1.2 × 10 ⁻³	0.34
4 × 10 ⁻⁴	2.5 × 10 ⁻⁴	1.5 × 10 ⁻³	0.37
5 × 10 ⁻⁴	1.4 × 10 ⁻³	2.2 × 10 ⁻³	0.37
0.001	5 × 10 ⁻³	2.3 × 10 ⁻³	0.44
0.001	2.5 × 10 ⁻³	2.1 × 10 ⁻³	0.33
0.001	1 × 10 ⁻³	1.9 × 10 ⁻³	0.34
0.001	5 × 10 ⁻⁴	1.4 × 10 ⁻³	0.37
0.001	3 × 10 ⁻⁴	1.2 × 10 ⁻³	0.32
0.001	2.5 × 10 ⁻⁴	1.2 × 10 ⁻³	0.32
0.001	7 × 10 ⁻⁵	4 × 10 ⁻⁴	0.36
0.001	7 × 10 ⁻⁵	4 × 10 ⁻⁴	0.36 ^c
0.001	5 × 10 ⁻⁴	1.4 × 10 ⁻³	0.37 ^c
0	5 × 10 ⁻⁴	2.4 × 10 ^{-3c}	

^a Added as tetrabutylammonium-cyanide.

^b The starting concentration of bis(pyridine)cobalt(III)hematoporphyrin-IX acetate was 2.7 × 10⁻⁵ M. In all the other runs it was 7 × 10⁻⁶ M.

^c Chloroform was dried by freshly prepared BDH A 4 molecular sieves lowering the water content from 0.01 M to less than 0.002 M [H₂O].

Acknowledgements. The support to this research by the Scientific Council of Croatia is gratefully acknowledged. The authors are grateful to Professor D. Grdenić and Drs N. Brničević and V. Jagodić for allowing the use of the facilities in their laboratories. We are indebted to the Fluka Comp. for obtaining the sample of tetrabutylammonium-cyanide.

REFERENCES

1. E. B. Fleischer and M. Krishnamurthy, *Ann. N. Y. Acad. Sci.* **206** (1973) 32.
2. P. Hambricht in K. M. Smith, (Ed) »Porphyrins and Metalloporphyrins«, Elsevier, Amsterdam, 1975, p. 268, and references therein.
3. F. R. Jensen and R. C. Kiskis, *J. Amer. Chem. Soc.* **97** (1975) 5820.
4. D. V. Stynes and B. R. James, *Chem. Comm.* (1973) 325.
5. R. J. McConnel, B. G. Overell, V. Petrow, and B. Sturgeon, *J. Pharm. Pharmacol.* **5** (1953) 179.
6. E. B. Fleischer, S. Jacobs, and L. Mestichelli, *J. Amer. Chem. Soc.* **90** (1968) 2527.

SAŽETAK**Kinetičko ispitivanje supstitucije piridina sa cijanidom u bis(piridin)kobalt(III)hematoporfirin-IX: Razlikovanje I_d i D mehanizama***M. Biruš i M. Pribanić*

Kinetičko ispitivanje reakcije cijanid-iona s bis(piridin)kobalt(III)hematoporfirin-IX kompleksom u kloroformu, pri čemu kao reakcijski produkt nastaje bis(cijanido)kobalt(III)hematoporfirin-IX kompleks, pokazuje da supstitucija aksijalno vezanih liganada slijedi čisti disocijacijski (D a ne I_d) mehanizam koji predviđa stvaranje (piridin)kobalt(III)hematoporfirin-IX pentakoordiniranog međuprodukta.

ZAVOD ZA KEMIJU, FARMACEUTSKO-
BIOKEMIJSKOG FAKULTETA SVEUČILIŠTA
U ZAGREBU
41000 ZAGREB

Prispjelo 11. kolovoza 1976.