

# First hydroxamic seconucleoside derivatives

---

Vrček, Valerije; Čaplar, Vesna; Uršić, Stanko

Source / Izvornik: **Croatica Chemica Acta, 1998, 71, 119 - 124**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:163:496056>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-07-14**



Repository / Repozitorij:

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



## First Hydroxamic Seconucleoside Derivatives

Valerije Vrček,<sup>a</sup> Vesna Čaplar,<sup>b</sup> and Stanko Uršić<sup>a</sup>

<sup>a</sup>Faculty of Pharmacy and Biochemistry, University of Zagreb, A. Kovačića 1,  
10001 Zagreb, Croatia (e-mail: stu@nana.pharma.hr)


<sup>b</sup>Laboratory of Supramolecular and Nucleoside Chemistry, Rugjer Bošković  
Institute, P.O.B. 1016, 10001 Zagreb, Croatia

Received May 21, 1997; revised September 15, 1997; accepted September 19, 1997

Nucleoside analogues possessing hydroxamate moiety were prepared from secouridine derivatives oxidized at C(2')-position. 3',5'-O-Isopropylidene-2',3'-secouridine was oxidized with CrO<sub>3</sub>/py complex to the carboxylate level and with DMSO/DCC to the aldehyde level. Both compounds (precursors) gave C(2')-nucleoside hydroxamic acids in the reaction with hydroxylamine or nitrosobenzene in the presence of Fe<sup>3+</sup> ions, respectively.

A number of hydroxamic acid analogues have been shown to inhibit DNA synthesis by inactivating the enzyme ribonucleotide reductase (RNR).<sup>1–4</sup> This metalloenzyme catalyzes the conversion of (ribo)nucleotides to deoxy(ribo)nucleotides and is therefore a potential target for the development of anticancer agents.<sup>5–8</sup> Hydroxamic acid moiety, R–CONHOH, is found to be the essential pharmacophore in the hydroxyurea, a clinically useful inhibitor of ribonucleotide reductase.<sup>9</sup> A variety of nucleoside analogues are also active as inhibitors of ribonucleotide reductase,<sup>10–12</sup> following the inhibition mechanism similar to that proposed for hydroxamates.<sup>13–15</sup> These results have initiated our interest in the design and synthesis of the nucleoside analogues incorporating hydroxamate moiety. A similar concept was devised by Farr *et al.*<sup>16</sup> Their compounds inhibited RNR activity, but were 10-fold less potent than hydroxyurea. We, therefore, rationalize that hydroxamate-nucleoside analogues which more closely resemble the RNR substrates should be prepared. Moreover, recent reports<sup>17–19</sup> that hydroxamate compounds increase the potency of nucleosides against HIV-1 give an additional importance to derivatives combining the structural fea-


tures of the above compounds. To prepare hydroxamic-nucleoside derivatives, we have used securidine-2',3'-dialdehyde as the starting material. We have already reported preliminary results on the reaction of nitrosobenzene with 2',3'-securidine dialdehyde giving hydroxamic-nucleoside product.<sup>20</sup> It is shown that this reaction does not proceed in the absence of ferric ion under the conditions employed. In the presented work, we also prepared monoaldehyde form **3** of 2',3'-securidine **1** (Scheme 1) in order to prevent the formation of cyclic hemiacetals of dialdehyde monohydrate.<sup>21,22</sup>



Scheme 1. a) acetone,  $H^+$ , RT, 2 h; b) DMSO/benzene/pyridine, TFA and DCC, RT, overnight; c)  $Fe^{3+}$ ,  $H^+$ , PhNO in acetone, RT, 6 h.

3',5'-*O*-Isopropylidene-2',3'-seconucleoside **2** was obtained according to the procedure described by Jones *et al.*<sup>23</sup> The oxidation of **2** to the aldehyde level was performed by dimethylsulfoxide (DMSO) and dicyclohexylcarbodiimide (DCC) in the presence of pyridinium trifluoroacetate (PTFA).<sup>24</sup> The aldehyde product **3** was obtained in 60% yield. In the NMR spectra of **3** the signals of aldehyde proton at 10.30 ppm and aldehyde carbon at 194.20 ppm were present, although the compound exists predominately as hydrate **3a**. The test with dinitrophenylhydrazine was positive.

In the reaction of aldehyde **3** with nitrosobenzene, we have observed in UV spectrum the appearance of the highly characteristic peak at 520 nm due to the formation of the iron(III)-mono-*N*-phenylhydroxamate complex **4**. The reaction was promoted both with proton and  $Fe^{3+}$  ions. Anyway, the presence of  $Fe^{3+}$  in the reaction mixture made it difficult to isolate a free hydroxamic-nucleoside conjugate from complex product **4**. This is due to the known property of hydroxamates to form a strong complex with  $Fe^{3+}$ .<sup>25,26</sup> To circumvent this difficulty, we have applied a different approach to the synthesis of hydroxamic-nucleoside derivatives (Scheme 2).




Scheme 2. a) CrO<sub>3</sub>/pyridine complex in CH<sub>2</sub>Cl<sub>2</sub>-DMF, AcOAc, *t*-butanol, RT, 20 h; b) NH<sub>2</sub>OH in MeOH, RT, 3 h; c) 40% AcOH, RT, 3 h.

One-step oxidation of primary alcohol **2** to the corresponding carboxylic *tert*-butyl ester **5**<sup>27</sup> was performed using chromium trioxide-pyridine complex<sup>28</sup> in dichloromethane-dimethylformamide (4:1) in the presence of *tert*-butanol and acetic anhydride (60% yield). 10% of the starting compound and 15% of acetylated by-product, 2'-acetyl-3',5'-*O*-isopropylidene-2',3'-secouridine (**6**), were also isolated from the reaction mixture. Omission of dimethylformamide from the reaction mixture led to poorer solubility of the oxidizing complex and consequently to slower oxidation. Therefore, in dichloromethane, the yield of oxidized product **5** was diminished (26%) and the ratio of acetylated product **6** doubled.

5'-*O*-Trityl-2',3'-secouridine **7** (Scheme 3.) could not be oxidized to *tert*-butyl carboxylates at C(2') and/or C(3') position by this method, presumably because of the complex cyclic hemiacetal structures of intermediary dialdehydes.<sup>21,22</sup> The only isolated products were the acetylated starting compound, 2',3'-di-*O*-acetyl-2',3'-secouridine (**8**), and 1-[2(*S*)-trityloxymethyl-3(*R,S*)-acetoxy-1,4-dioxan-6(*R*)-yl]uracil (**9**). 1,4-Dioxacyclohexane structure **9** is a diastereomeric mixture resulting from acetylation of the previously formed intramolecular aldehyde hemiacetal. Also, an attempt at oxidation of 5'-*O*-trityl-2',3'-secouridine dialdehyde **10** resulted only in a complex mixture of diastereoisomeric cyclic acetals.

The method used for the preparation of 2'-hydroxamate derivative **11** (Scheme 2.) is similar to that described for the synthesis of the first sugar-



C(2') and/or C(3') *tert*-butyl carboxylates

Scheme 3. a) CrO<sub>3</sub>/pyridine complex in CH<sub>2</sub>Cl<sub>2</sub>-DMF, AcOAc, *t*-butanol, RT, 20 h.

hydroxamic acid.<sup>29</sup> Solution of sodium methoxide was added to hydroxylamine hydrochloride to prepare a free hydroxylamine. After removing sodium chloride by filtration hydroxylamine was reacted with *tert*-butyl-3',5'-*O*-isopropylidene-2',3'-secouridine-2'-carboxylate **5** giving 3',5'-*O*-isopropylidene-2',3'-secouridine-2'-hydroxamic acid **11** (65% yield). Deprotection was achieved in 40% acetic acid for 3 hours at room temperature giving hydroxamic-nucleoside derivative **12**<sup>30</sup> (65% yield). As far as we know, it is the first seconucleoside analogue incorporating hydroxamic moiety. We are continuing investigations on hydroxamic-nucleoside analogues with other nucleobases. Metal coordination properties and biological testing of compounds described will be reported in due course.

## REFERENCES AND NOTES

1. J. C. Swarts, M. A. S Aquino, J. Y. Han, K. Y. Lam, and A. G. Sykes, *Biochim. Biophys. Acta Protein Struct. Mol. Enzymol.* **1247** (1995) 215–224.
2. I. K. Larsen, B. -M. Sjöberg, and L. Thelander, *Eur. J. Biochem.* **125** (1982) 75–81.
3. B. van't Riet, G. L. Wampler, and H. L. Elford, *J. Med. Chem.* **22** (1979) 589–591.
4. H. L. Elford, G. L. Wampler, and B. van't Riet, *Cancer Res.* **39** (1979) 844–851.
5. P. Reichard, *Science* **260** (1993) 1773–1777.
6. J. Stubbe, *Adv. Enzymol. Relat. Areas Mol. Biol.* **63** (1990) 349–374.
7. J. Stubbe and W. A. van der Donk, *Chem. Biol.* **2** (1995) 793–801.
8. M. N. Prichard and C. Shipman, *Chemotherapy* **41** (1995) 384–395.
9. S. P. Gupta, *Chem. Rev.* **94** (1994) 1507–1551.
10. K. C. Xie and W. Plunkett, *Cancer Res.* **56** (1996) 3030–3037.
11. T. E. Lehmann and A. Berkessel, *J. Org. Chem.* **62** (1997) 302–309.
12. S. Salowe, J. M. Bolliger, M. Ator, J. Stubbe, J. McCracken, J. Peisach, M. C. Samano, and M. J. Robins, *Biochemistry* **32** (1993) 12749–12760.
13. M. J. Robins, Z. Guo, and S. F. Wnuk, *J. Am. Chem. Soc.* **119** (1997) 3637–3638.
14. R. Eliasson, E. Pontis, F. Eckstein, and P. Reichard, *J. Biol. Chem.* **265** (1994) 26116–26120.

15. R. Ingemarson and L. Thelander, *Biochemistry* **35** (1996) 8603–8609.
16. R. A. Farr, P. Bey, P. S. Sunkara, and B. J. Lippert, *J. Med. Chem.* **32** (1989) 1879–1885.
17. S. D. Malley, J. M. Grange, F. Hamedi-Sangsari, and J. R. Vila, *Proc. Natl. Acad. Sci. USA* **91** (1994) 11017–11021.
18. F. Lori, A. Malykh, A. Cara, D. Sun, J. N. Weinstein, J. Lisiewicz, and R. C. Gallo, *Science* **266** (1994) 801–805.
19. W. -Y. Gao, H. Mitsuya, J. S. Driscoll, and D. G. Johns, *Biochem. Pharmacol.* **50** (1995) 274–276.
20. S. Uršić, B. Nigović, V. Vrček, and V. Pilepić, *Tetrahedron Lett.* **36** (1995) 9547–9550.
21. O. Howarth, A. S. Jones, R. T. Walker, and P. G. Wyatt, *J. Chem. Soc. Perkin Trans. II* (1984) 261–265.
22. J. P. Neenan, S. M. Opitz, C. L. Cooke, M. A. Ussery, T. C. Morrill, and L. M. Eckel, *Bioorg. Med. Chem. Lett.* **6** (1996) 1381–1386.
23. A. S. Jones, M. J. McClean, H. Tanaka, R. T. Walker, J. Balzarini, and E. de Clercq, *Tetrahedron* **41** (1985) 5965–5972.
24. K. E. Pfitzner and J. G. Moffatt, *J. Am. Chem. Soc.* **87** (1965) 5670–5678.
25. B. F. Matzanke, G. Mueller-Matzanke, and K. N. Raymond, *Siderophore Mediated Iron Transport; Chemistry, Biology and Physical Properties*, in: T. M. Loehr, H. B. Gray, and A. B. Lever (Eds.), *Physical Bioorganic Chemistry*, VCH Publishers, New York, 1989.
26. M. J. Miller and F. Malouin, *Acc. Chem. Res.* **25** (1993) 241–252.
27. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ/ppm: 9.0 (bs, 1H, NH), 7.51 (d, 1H, *J* = 8.13 Hz, 6-H), 6.10 (s, 1H, 1'-H), 5.82 (d, 1H, *J* = 8.14 Hz, 5-H), 4.09–3.70 (m, 5H, 3'-H<sub>2</sub>, 4'-H, 5'-H<sub>2</sub>), 1.49 (s, 9H, *t*-Bu), 1.42 and 1.41 (2s, 2x3H, *i*Pr). <sup>13</sup>C-NMR (CDCl<sub>3</sub>) δ/ppm: 164.80 (C-2'), 163.42 (C-4), 150.66 (C-2), 140.52 (C-6), 102.82 (C-5), 98.31 (CMe<sub>2</sub>), 84.24 (CMe<sub>3</sub>), 79.94 (C-1'), 71.60 (C-4'), 61.76 and 61.70 (C-3' and C-5'), 27.51 (CMe<sub>3</sub>), 24.16 and 22.20 (CMe<sub>2</sub>). *Anal.* Calcd. for C<sub>16</sub>H<sub>24</sub>N<sub>2</sub>O<sub>7</sub> (*M<sub>r</sub>* = 356.38): C 53.93, H 6.79, N 7.86%; found: C 53.98, H 7.03, N 7.71%.
28. E. J. Corey and B. Samuelsson, *J. Org. Chem.* **49** (1984) 4735–4735.
29. S. Uršić, B. Zorc, V. Pilepić, and D. Vikić-Topić, *Croat. Chem. Acta* **65** (1992) 851–857.
30. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ/ppm: 7.52 (d, 1H, *J* = 8.0 Hz, 6-H), 6.04 (s, 1H, 1'-H), 5.61 (d, 1H, *J* = 8.0 Hz, 5-H), 4.0–4.5 (m, NHOH, 2x OH), 3.55–3.33 (m, 5H, 3'-H<sub>2</sub>, 4'-H, 5'-H<sub>2</sub>). <sup>13</sup>C-NMR (DMSO-d<sub>6</sub>) δ/ppm: 163.73 (C-2'), 161.65 (C-4), 151.29 (C-2), 142.48 (C-6), 101.49 (C-5), 81.92 and 81.40 (C-4' and C-1'), 60.77 and 60.58 (C-3' and C-5'). *Anal.* Calcd. for C<sub>9</sub>H<sub>13</sub>N<sub>3</sub>O<sub>7</sub> (*M<sub>r</sub>* = 275.22): C 39.28, H 4.76, N 15.27%; found: C 39.33, H 4.75, N 15.30%.

**SAŽETAK****Prvi derivati hidroksamskih sekonukleozida**

*Valerije Vrček, Vesna Čaplar i Stanko Uršić*

Nukleozidni derivati s hidroksamskom skupinom pripremljeni su iz sekouridina oksidiranog na položaju C(2'). 3',5'-*O*-Izopropiliden-2',3'-sekouridin oksidiran je s pomoću kompleksa CrO<sub>3</sub>/piridin do karboksilata, a pomoću DMSO/DCC do aldehida. Karboksilatni derivat u reakciji s hidroksilaminom, odnosno aldehydni derivat u reakciji s nitrozobenzenom u prisutnosti iona Fe<sup>3+</sup> daje C(2')-hidroksamski nukleozid.