

# Formation of hydroxamic acids promoted by metal ions. Interaction of nitroso group with coordinated carbonyl group

---

**Radivoj, Denis; Pilepić, Viktor; Uršić, Stanko**

*Source / Izvornik:* **Croatica Chemica Acta, 1996, 69, 1633 - 1638**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

*Permanent link / Trajna poveznica:* <https://urn.nsk.hr/urn:nbn:hr:163:853363>

*Rights / Prava:* [In copyright](#)/[Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2024-07-14**



*Repository / Repozitorij:*

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



## Formation of Hydroxamic Acids Promoted by Metal Ions. Interaction of Nitroso Group with Coordinated Carbonyl Group\*

*Denis Radivoj,<sup>a</sup> Viktor Pilepić,<sup>b</sup> and Stanko Uršić<sup>b,\*\*</sup>*

*<sup>a</sup>Air Force Base Pleso, Croatia*

*<sup>b</sup>Faculty of Pharmacy and Biochemistry, University of Zagreb,  
A. Kovačića 1, Zagreb, Croatia*

Received September 1, 1996; revised October 5, 1996; accepted October 11, 1996

Hydroxamic acids are formed in the interaction of nitroso group of substituted nitrosobenzenes with carbonyl group of glyoxylate coordinated on iron(III) ion.

Interaction of nitroso compounds having a nitroso group on carbon (C-nitroso group) with carbonyl group of  $\alpha$ -oxo acids<sup>1</sup> and aldehydes<sup>1a,b,2</sup> leads to the formation of hydroxamic acids, a class of compounds of exceptional biochemical, pharmaceutical and industrial importance.<sup>3–5</sup> For example, the roles of hydroxamic acids as enzyme inhibitors<sup>4</sup> or neoplastic cell differentiation inducers<sup>5</sup> are currently investigated. Nitroso group is known to act as an electrophile in many of its reactions,<sup>6</sup> but it also possesses nucleophilic properties. This fact, predicted also by theoretical studies,<sup>7</sup> was demonstrated by both solution and solid-state NMR spectroscopy,<sup>8,9</sup> as well as by kinetic studies of its nucleophilic reactions.<sup>1,2</sup> These include, among others, interactions of substituted nitrosobenzenes and aliphatic nitroso compound 2-methyl-2-nitrosopropane with biochemically important carbonyl substrates


---

\* Dedicated to the memory of Professor Stanko Borčić.

\*\* Author to whom correspondence should be addressed.

such as glyoxylate,<sup>1a,c</sup> pyruvic acid,<sup>1b</sup> glyoxylic acid,<sup>1a</sup> and acetaldehyde.<sup>1b</sup> Products of these reactions are, as mentioned, hydroxamic acids. The reactions are complex, and nitroso group acts as a nucleophile in the first reaction step, where nucleophilic attack of nitroso nitrogen on carbonyl carbon results in the C–N bond formation, leading to the corresponding dipolar addition intermediate.<sup>2a</sup> To our knowledge, there is no evidence reported that nitroso group can interact in analogous manner with carbonyl group coordinated on a metal ion.

In this work, we report that substituted nitrosobenzenes react with glyoxylate coordinated on iron(III) ion, giving the corresponding *N*-phenyl hydroxamic acids. The reaction involves the formation of the complex intermediate (1), Scheme 1, resulting from the interaction of nitroso nucleophile



with  $\alpha$ -oxo group of glyoxylate coordinated on iron(III) ion, thus catalyzing the subsequent formation of the nitrosocarbinolic intermediate (2) which leads to the product, hydroxamic acid (3). The evidence includes:

*i*) The observation that the reaction product of the reaction of nitroso nucleophile with glyoxylic carbonyl group, under the conditions where the rate of the reaction *via* uncomplexed glyoxylate/glyoxylic acid was kinetically insignificant in the overall process, was the corresponding hydroxamic acid.

*ii*) The observation of the metal ion catalysis (Figure 1). The catalysis is rationalized by the obtained linear dependence of the pseudo-first order rate constants for the formation of hydroxamic acid on the concentration of mono glyoxylato iron(III) complex, Figure 2, expressed by  $k_{\text{obs}} = k_0 + k_1 [\text{FeL}^{2+}]$  ( $\text{L}^- = \text{glyoxylate}$ ), where  $k_0$  refers to the ferric ion uncatalysed reaction path, as described in detail earlier.<sup>1,2</sup> Values of  $k_0$ <sup>13</sup> and  $k_1$  are  $0.0067 \text{ s}^{-1}$  and  $0.516 \text{ s}^{-1} \text{ M}^{-1}$  respectively, under the conditions employed.

*iii*) The slope of the straight line of  $\log k_{\text{obs}}$  *vs.* Hammett  $\sigma^+$  parameters, obtained for the reaction of glyoxylate with substituted nitrosobenzenes in the presence of ferric ions differs substantially from that of the corresponding line of  $\log k_{\text{obs}}$  *vs.* Hammett  $\sigma^+$  parameters in the absence of iron(III) ions (Figure 3). The negative value of this slope in the case of reaction of




Figure 1. Dependence of the pseudo-first order rate constants for the formation of *N*-phenylformhydroxamic acid in the reaction of glyoxylic acid/glyoxylate with nitrosobenzene on the iron(III) ion concentration at pH = 1.56. Individual rate constants are determined using the methods reported.<sup>1a,b;2a,c</sup> In water, at 25 °C, ionic strength: 2.0 M (NaClO<sub>4</sub>/Fe(ClO<sub>4</sub>)<sub>3</sub>), total glyoxylate /glyoxylic acid: 0.05 M.<sup>10,13</sup>




Figure 2. Dependence of the pseudo-first order rate constants for the formation of *N*-phenylformhydroxamic acid in the reaction of glyoxylic acid/glyoxylate with nitrosobenzene on the monoglyoxylato iron(III) complex concentration.<sup>13</sup> At pH = 1.56, in water, at 25 °C, ionic strength: 2.0 M (NaClO<sub>4</sub>/Fe(ClO<sub>4</sub>)<sub>3</sub>), total glyoxylate/glyoxylic acid: 0.05 M. Rate constants were determined as described earlier.<sup>1a,b;2a,c</sup> Fraction of FeL<sup>2+</sup> complex was calculated (L<sup>-</sup> = glyoxylate) using an equilibrium quotient  $K_{eq} = 7.76$ , defined as  $FeL^{2+} \cdot H^+ / Fe^{3+} \cdot HL$  for the formation of monoglyoxylato iron(III) complex, as determined spectrophotometrically, at pH = 1.56, (under the conditions employed in kinetic measurements, by measuring the increase of absorbance at 450 nm due to the formation of the complex).




Figure 3. Plot of the  $k_{\text{obs}}$  vs. Hammett  $\sigma^+$  parameters for the reactions of glyoxylate with the substituted nitrosobenzenes R-Ph-NO. In the presence of 0.009 M  $\text{Fe}^{3+}$ , (total glyoxylate = 0.194 M, pH = 1.55, ionic strength = 0.2 M),  $\rho^+ = -1.22$ ,  $r = 0.970$ , (open circles). In the absence of  $\text{Fe}^{3+}$ , (total glyoxylate = 0.194 M, pH = 1.55, ionic strength = 0.2 M),  $\rho^+ = -2.20$ ,  $r = 0.998$ , (solid circles).

glyoxylate/glyoxylic acid with substituted nitrosobenzenes, as well as in similar reactions,<sup>1,2</sup> was related to the nucleophilic attack of nitroso nucleophile in the first reaction step. Therefore, it seems reasonable to conclude from the evidence obtained that the metal ion intervenes in this reaction step.

iv) The observed solvent deuterium isotope effects  $k_{\text{D}_2\text{O}}/k_{\text{H}_2\text{O}}$  (uncorrected) were as follows: 1.11 (0.07) at  $\text{Fe}^{3+} = 0.073$  M, total glyoxylate = 0.01 M, pH = 1.15, pD(reading) = 0.80, in 95%  $\text{D}_2\text{O}$ ; 1.19 (0.08) at  $\text{Fe}^{3+} = 0.073$  M, total glyoxylate = 0.05 M, pH = 1.32, pD(reading) = 1.03, in 95%  $\text{D}_2\text{O}$ ; 1.01 (0.01)<sup>14</sup> at  $\text{Fe}^{3+} = 0.008$  M, total glyoxylate = 0.084 M, pH = 1.32, pD(reading) = 1.02, in 84%  $\text{D}_2\text{O}$ . These isotope effects are small and inverse. This is consistent with the preequilibrium protonation<sup>15,16</sup> of (1) which leads to the nitrosocarbonylic polar intermediate (2). Such a preequilibrium protonation could lead to the inverse solvent deuterium isotope effect ( $k_{\text{D}_2\text{O}}/k_{\text{H}_2\text{O}}$ ) of a magnitude of 2–4.<sup>15,16</sup> However, the observed small inverse solvent deuterium isotope effects should result from the mutual cancellation of the corresponding solvent isotope effect on iron(III) complexation<sup>17</sup> with glyoxylate (under the conditions employed) and the solvent isotope effect on preequilibrium protonation of complex intermediate (1).

It should be noted that C-nitroso compounds could be of interest as antiretroviral<sup>18</sup> and cytotoxic<sup>19</sup> agents. Hence, a better insight into its reactivity toward the coordinated carbonyl compounds of biochemical importance could be helpful in C-nitroso drug design.

*Acknowledgments.* – We thank Miss Jelena Bužančić for collaboration and the Croatian Research Council for support.

## REFERENCES

1. a) V. Pilepić and S. Uršić, *Tetrahedron Lett.* **35** (1994) 7425. b) S. Uršić, V. Pilepić, V. Vrček, M. Gabričević, and B. Zorc, *J. Chem. Soc., Perkin Trans. 2* (1993) 509. c) M. D. Corbett and B. R. Corbett, *J. Org. Chem.* **45** (1980) 2834.
2. a) S. Uršić, *Helv. Chim. Acta* **76** (1993) 131. b) O. Kronja, J. Matijević-Sosa, and S. Uršić, *J. Chem. Soc., Chem. Commun.* (1987) 463. c) S. Uršić, B. Nigović, V. Vrček, and V. Pilepić, *Tetrahedron Lett.* **36** (1995) 9547.
3. a) B. F. Matzanke, G. Mueler-Matzanke, and K. N. Raymond, *Siderophore Mediated Iron Transport; Chemistry, Biology and Physical Properties*, in: T. M. Loehr, H. B. Gray, and A. B. P. Lever (Eds.), *Physical Bioorganic Chemistry*, VCH Publishers, New York, 1989. b) C. Hider and A. D. Hall, in: R. W. Hay, J. R. Dilworth, and K. B. Nolan (Eds.) *Perspectives in Bioinorganic Chemistry*, JAI Press, London, 1991, p. 209.
4. a) B. Walcheck, J. Kahn, J. M. Fisher, B. B. Wang, R. S. Fisk, D. G. Payan, C. Feehan, R. Betageri, K. Darlak, A. F. Spatola, and T. K. Kishimoto, *Nature* **380** (1996) 720. b) R. R. L. Hamer, J. J. Tegeler, E. S. Kurtz, R. C. Allen, S. C. Bailey, M. E. Elliott, L. Hellyer, G. C. Helsley, P. Przekop, B. S. Freed, J. White, and L. L. Martin, *J. Med. Chem.* **39** (1996) 246.
5. V. M. Richon, Y. Webb, R. Merger, T. Sheppard, B. Jursic, L. Ngo, F. Civoli, R. Breslow, R. A. Rifkind, and P. A. Marks, *Proc. Natl. Sci. U.S.A.* **93** (1996) 5705.
6. P. Zuman and B. Shah, *Chem. Rev.* (1994) 1621.
7. P. Politzer and R. Bar-Adon, *J. Phys. Chem.* **91** (1987) 2069.
8. K. G. Orrell, V. Šik, and D. Stephenson, *Magn. Reson. Chem.* **25** (1987) 1007.
9. M. D. Lumsden, G. Wu, R. E. Wasylishen, and R. D. Curtis, *J. Am. Chem. Soc.* **115** (1993) 2825.
10. Glyoxylate and glyoxylic acid are present in solution mainly in the form of hydrated species<sup>11</sup> and glyoxylic acid ( $pK_a \approx 2$ , for the dehydrated glyoxylic acid, see for example Ref. 12) is much more hydrated than glyoxylate itself.
11. J. E. Meany and Y. Pocker, *J. Am. Chem. Soc.* **113** (1991) 6155, and references therein.
12. P. E. Sørensen, K. Bruhn, and F. Lindeløv, *Acta Chem. Scand. A*, **28** (1974) 162.
13. Under the conditions employed, the uncatalyzed reaction does not contribute significantly to the overall rate of the reaction.  $k_o$  of  $0.0067 \text{ s}^{-1}$  refers to the total glyoxylate concentration of 1 M.
14. In the case of this solvent isotope effect, there is about 30% of uncatalyzed reaction. Solvent deuterium isotope effect on  $k_o$  is  $k_{D_2O}/k_{H_2O} = 1.92$ , at  $pH = 1.24$ . Therefore, cancellation of the inverse isotope effect could be partly a consequence of the normal isotope effect in uncatalyzed reaction.
15. P. M. Laughton and R. E. Robertson, in: J. F. Coetzee and C. D. Ritchie (Eds.), *Solute-Solvent Interactions*, Marcel Dekker, New York, 1969, p.399.
16. See also discussion in Ref. 1b and 2a.

17. Obviously, normal solvent deuterium isotope effect on the equilibrium of the complex formation of  $\text{FeL}^{2+}$  should be expected. It is well known that  $\Delta pK_a$  between  $\text{D}_2\text{O}$  and  $\text{H}_2\text{O}$  for most weak acids lies between 0.3 and 0.6.<sup>15</sup>
18. W. G. Rice, C. A. Schaeffer, L. Graham, M. Bu, J. S. McDougal, S. L. Orloff, F. Villinger, M. Young, S. Oroszlan, M. R. Fesen, Y. Pommier, J. Mendeleyev, and E. Kun, *Nature* **361** (1993) 473.
19. J. Mendeleyev, E. Kirsten, A. Hakam, K. G. Buki, and E. Kun, *Biochem. Pharmacology* **50** (1995) 705.

### SAŽETAK

#### **Nastajanje hidroksamskih kiselina potpomognuto kovinskim ionima. Međudjelovanje nitrozo-skupine i koordinirane karbonilne skupine**

*Denis Radivoj, Viktor Pilepić i Stanko Uršić*

Međudjelovanjem karbonilne skupine glioksilata koordinirane na ion  $\text{Fe(III)}$  s nitrozo-skupinom supstituiranih nitrozobenzena nastaju odgovarajuće *N*-fenil supstituirane hidroksamske kiseline.