

# Metilensko modrilo - lijek, bojilo, indikator

---

**Zorc, Branka; Pavić, Kristina**

*Source / Izvornik:* **Farmaceutski glasnik, 2018, 73, 99 - 108**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

*Permanent link / Trajna poveznica:* <https://um.nsk.hr/um:nbn:hr:163:568787>

*Rights / Prava:* [In copyright](#) / [Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2024-07-19**



*Repository / Repozitorij:*

[Repository of Faculty of Pharmacy and Biochemistry University of Zagreb](#)



# Metilensko modrilo – lijek, bojilo, indikator


BRANKA ZORC, KRISTINA PAVIĆ

Sveučilište u Zagrebu, Farmaceutsko-biokemijski fakultet,  
Zavod za farmaceutsku kemiju, A. Kovačića 1, 10000 Zagreb

Metilensko modrilo ili metiltioninijev klorid naziv je za 3,7-bis(dimetilamino)-fenotiazin-5-ijev klorid (slika 1.).

Zajedno s azurom A, B i C pripada fenotiazin-skim bojilima (razlikuju se po stupnju metilacije amino skupina). Prvi ga je sintetizirao Heinrich Caro 1876. godine. Od tada se koristi kao lijek, bojilo i indikator. Monografija o metilenskom modrilu nalazi se u većini farmakopeja, između ostalog u europskoj i američkoj farmakopeji (1, 2). U USP 37 su osim metilenskog modrila oficinalna i dva pripravka: Injekcija metilenskog modrila (slika 2.) i Injekcija metilenskog modrila za veterinarsku upotrebu. Metilensko modrilo se nalazi na listi esencijalnih lijekova Svjetske zdravstvene organizacije u skupini antidota i ostalih supstancija koje se koriste kod trovanja (3).

Metilensko modrilo je prvi potpuno sintetski spoj koji se koristio kao lijek. Paul Guttman i Paul Erlich uveli su ga u terapiju malarije još 1891. godine. Do tada se kao antimalarik koristio kinin. Kinin se izolirao iz kore južnoameričke biljke kininovca (*Cinchona officinalis* L.), što je znatno ograničavalo opskrbu lijekom. Upravo je otkriće da jedna u potpunosti sintetska tvar, metilensko modrilo, djeluje protiv malarije, omogućilo bolje provođenje antimalarijske terapije (4). U Drugom Svjetskom ratu američki vojnici u tropskim dijelovima Pacifika dobivali su metilensko modrilo za prevenciju i liječenje malarije. Zbog bojanja urina i bjeloočnica, vojnici su ga uzimali nerado pa se ta praksa ukinula. No, u novije vrijeme interes za metilenskim modrilom kao antimalarikom ponovo raste (6–10). Sada je u tijeku nekoliko velikih kliničkih studija u kojima se ispituje antimalarijski učinak metilenskog modrila (11–13).



Slika 1. Strukturna formula metilenskog modrila.




Slika 2. Otopina metilenskog modrila za intravensku primjenu.

Danas se intenzivno proučava mogući utjecaj metilenskog modrila na razvoj Alzheimerove bolesti (14, 15). Naime, metilensko modrilo u *in vitro* uvjetima i na animalnim modelima inhibira agregaciju tau proteina i utječe na disocijaciju amiloida pa je pod zaštićenim imenima, prvo kao Rember, zatim pod nazivom LMTX ili TRx0237 ušao u klinička ispitivanja (trenutno je TRx0237 u trećoj fazi) kao potencijalni lijek za Alzheimerovu bolest (16). Najnovija istraživanja pokazuju da metilensko modrilo u malim dozama ( $0,5\text{--}4\text{ mg kg}^{-1}$ ) poboljšava kratkotrajnu memoriju u ljudi (17), a *in vitro* istraživanja dodatno pokazuju učinke na usporavanje starenja stanica i sprječavanje disfunkcije mitohondrija, osobito enzima kompleks IV (18, 19). Također je dokazano njegovo neuroprotektivno djelovanje u eksperimentalnim modelima ishemičkog moždanog udara i Parkinsonove bolesti (20). Osim toga, metilensko modrilo modulira agregaciju intermedijera huntingtina pa pokazuje protektivni učinak u modelima još jedne neurodegenerativne bolesti – Huntingtonove bolesti (21).

Na američkim tržištu registrirano je nekoliko lijekova koji sadrže metilensko modrilo, a upotrebljavaju se u terapiji infekcija urinarnog trakta (22). Ti lijekovi sadrže kombinaciju antiseptika (metilensko modrilo, urotropin, benzojeva kiselina), analgetika i antiinflamatorika (fenil-salicilat) i spazmolitika (hiosciamin, atropin ili kombinacije alkaloida velebilja) (23).

Metilensko modrilo se može upotrijebiti kao antidot kod trovanja cijanidima, ugljikovim monoksidom (24), nitratima i nitritima (25), ali se u tu svrhu više ne upotrebljava. U farmakološkim dozama ( $1\text{--}2\text{ mg kg}^{-1}$ ) djeluje kao reducens, a u višim dozama kao antioksidans. Upravo zbog reduktivnih svojstava koristi se u terapiji methemoglobinemije (MhB) uzrokovane nekim lijekovima ili toksinima (slika 3.). U našem organizmu methemoglobin se reducira u hemoglobin pomoću nekoliko methemoglobin reduktaza koje koriste NADH ili NADPH kao koenzime. Ako nastanu velike količine methemoglobina, kapaciteti enzima su nedovoljni pa je potrebno koristiti metilensko modrilo kao antidot (22). No, velike doze ( $20\text{--}30\text{ mg kg}^{-1}$ ) metilenskog modrila induciraju methemoglobinemiju (26).

Metilensko modrilo inhibira enzim gvanilat ciklazu i monoamino oksidazu (MAO). Inhibicija prvog enzima sprječava nakupljanje cikličkog gvanozin monofosfata (cGMP)



Slika 3. Redukcija methemoglobina u prisutnosti metilenskog modrila.

zbog čega se smanjuje osjetljivost krvnih žila na cGMP-ovisne vazodilatatore kao što su dušikov(II) oksid i ugljikov(II) oksid (27). Zbog toga se metilensko modriilo može koristiti pri ekstremnoj hipotenziji uzrokovanoj sepsom (28) ili tijekom kirurških zahvata na srcu pri ugradnji prenosnica (29). Inhibicija drugog enzima dovodi do serotoniniskog sindroma ako se metilensko modriilo koristi intravenski u dozama većim od 5 mg/kg u kombinaciji sa selektivnim i neselektivnim inhibitorima ponovne pohrane serotonina (lijevovi iz skupine antidepresiva i anoreksika) (30). Zbog toga je kontraindicirana upotreba metilenskog modrila s tim lijekovima. Ne smije se koristiti uz MAO inhibitore (npr. uz antidepresiv tranilcipromin i moklobemid ili uz anti-parkinsonike selegilin i razagilin).


Nekoliko je studija pokazalo da metilensko modriilo može biti korisno u profilaksi i smanjenju encefalopatije uzrokovane ifosfamidom. Preporuča se njegova upotreba dan prije početka terapije ifosfamidom i tri puta dnevno tijekom provođenja kemoterapije (31) ili svakih 6 sati (32).

Virucidno djelovanje metilenskog modrila i drugih fenotiazinskih bojila u kombinaciji sa svjetlošću poznato je više od 70 godina (33). Takve kombinacije našle su primjenu u terapiji psorijaze (34), Kaposijevog sarkoma (35), infekcija uzrokovanih rezistentnim sojevima *Staphylococcus aureus* (36), virusom zapadnog Nila (37), HIV-om (38) te virusima koji uzrokuju hepatitis B i C (39, 40). No, kombinacija metilenskog modrila sa svjetlošću može uzrokovati oštećenje DNA i razvoj karcinoma (41, 42). Od ostalih nuspojava treba spomenuti hipertenziju, vrtoglavicu, zbunjenost, glavobolju, vrućicu, mučninu, povraćanje, bolove u abdomenu, bojanje kože i bjeloočnica, promjenu boje urina i fekalija (43, 44). U osoba koje su deficitentne na glukoza-6-fosfat dehidrogenazu (deficijencija G6PD, favizam) metilensko modriilo (kao i neki drugi lijekovi, npr. klorokin, primakin, sulfonamidi) uzrokuje hemolitičku anemiju (22). Naime, taj je enzim uključen u put pentozna-fosfata koji je posebno važan u crvenim krvnim stanicama. U literaturi je opisana i upotreba metilenskog modrila kao pomoćnog sredstva u terapiji vazoplegičnog sindroma nakon kirurškog zahvata na srcu (45, 46). Nekoliko pretkliničkih istraživanja ukazuju na selektivnu indukciju apoptoze stanica karcinoma nakon tretiranja metilenskim modrilom i srodnim fenotiazinskim bojilima (47, 48). Ispitivano je i djelovanje kombinacije metilenskog modrila i aoksina u fotodinamskoj terapiji karcinoma (49).

Zanimljiva je i upotreba metilenskog modrila kao placeba. Pacijentima je sugerirano da je plava boja urina znak učinkovitosti terapije. U psihijatriji se pomoću metilenskog modrila kontrolirala suradljivost pacijenata te prihvaćanje režima liječenja. Zbog toga ne začuđuje činjenica da je upravo metilensko modriilo bilo vodeći spoj iz kojeg se razvio prvi neuroleptik klorpromazin (50).


U biologiji se metilensko modriilo upotrebljava u nekoliko različitih postupaka za bojanje nukleinskih kiselina (bojanje po Wrightu, bojanje po Jenneru) (51). Koristi se i kao indikator preživljavanja stanica u testovima *in vitro*: žive stanice se ne oboje

jer se u njima metilensko modriilo reducira u bezbojnu formu, a mrtve stanice ostaju obojene (52). Važna je i njegova primjena u kirurgiji za označavanje tkiva na kojem će se obaviti manji kirurški zahvat (npr. odstranjivanje polipa) (53), za vizualizaciju limfnih putova pri odstranjivanju limfnih čvorova (slika 4.) (54) te u kemoendoskopiji za identificiranje displazije i prekanceroznih lezija u gastrointestinalnom traktu (55). U ortopedijskim operacijama dodaje se u koštani cement što omogućava lakše razlikovanje cementa od nativne kosti. Dodatno, ubrzava stvrđavanje koštanog cementa (56). Metilensko modriilo specifično boji živčana vlakna pa se na taj način proučavala inervacija mišića, kože i unutarnjih organa (slika 4.) (57).



**Slika 4.** Primjena metilenskog modrila za vizualizaciju limfnih čvorova (58).

Metilensko modriilo koristi se kao redoks indikator u analitičkoj kemiji (59): oksidirana forma metilenskog modrila je plava, a reducirana forma bezbojna (slika 5.). Nadalje, koristi se za fotometrijsko određivanje sulfida (60). Na tržištu je dostupan kit pod nazivom *Spectroquant Sulfide Test* za rutinsku analizu sulfida koji je npr.



**Slika 5.** Oksidirana i reducirana forma metilenskog modrila.

pogodan za brzu detekciju bakterija koje reduciraju sulfate (61). Metilensko modriilo se koristi i u tzv. MBAS testu (*Methylene Blue Active Substances Assay*), kojim se dokazuje prisutnost anionskih surfaktanata u vodi (62). Metilenskim testom može se odrediti i kapacitet gline za apsorpciju kationa iz otopine. Za tu svrhu proizveden je kit pod nazivom *Methylene Blue Kit* (63). U sintetskoj organskoj kemiji metilensko modriilo može se koristiti za sintezu peroksida. Reakcija ide preko singletnog kisika koji nastaje reakcijom metilenskog modriila s kisikom u prisutnosti svjetlosti (64).

Metilensko modriilo ima i komercijalnu primjenu: koristi se u akvarijima za tretman i prevenciju infekcija gljivicama i bakterijama (65) te u tekstilnoj industriji kao boja za tkanine.

Sumarno, na slici 6 prikazana su područja u kojima se upotrebljavao, upotrebljava ili će se moguće upotrebljavati metilensko modriilo.



Slika 6. Prijašnja, sadašnja i potencijalna upotreba metilenskog modriila.

# Methylene blue - medicine, dye, indicator

B. Zorc, K. Pavić

## Abstract

Methylene blue is a trivial name of 3,7-bis(dimethylamino)-phenothiazine-5-ium chloride). It has many uses in medicine, pharmacy, biology and chemistry. It is on the World Health Organization's List of Essential Medicines. From the end of 19th century to the end of Second World War methylene blue was used in therapy and prevention of malaria. Interest in its use as an antimalarial agent has recently been revived and several clinical trials are in progress. Another clinical trials investigate usefulness of methylene chloride for treatment of Alzheimer's dementia, ischemic stroke, Parkinson and Huntington's diseases.

Several papers report that methylene blue and related phenothiazine dyes appear to induce selective cancer cell apoptosis. Combined with light it has virucidal properties as well and combined with light and auxin it has a potential for the photodynamic treatment of cancer. Intravenous injections of methylene blue are used for the treatment of methemoglobinemia and as antidote in cyanide, carbon monoxide, nitrate and nitrite poisoning. It is used before initiation and during ifosfamide chemotherapy for treatment and prevention of ifosfamide neurotoxicity. In combination with other drugs it is used in treatment of urinary infection.

The most important side effects of methylene blue are hypertension, dizziness, headache, nausea, staining of skin and coloration of urine. Taken together with MAO inhibitors may precipitate serotonin syndrome. It causes hemolytic anemia in carriers of the glucose-6-phosphate dehydrogenase deficiency.

In surgery, endoscopy and biology methylene blue is used as a dye in different staining procedures and in analytical chemistry as a redox indicator, in sulfide analysis and in detection of anionic surfactants.

1. European Pharmacopoeia. 8<sup>th</sup> ed. Strasbourg: Council of Europe, 2015.
2. USP 37 – NF 32. Baltimore: United Book Press, Inc., 2013.
3. [http://apps.who.int/iris/bitstream/10665/93142/1/EML\\_18\\_eng.pdf?ua=1/](http://apps.who.int/iris/bitstream/10665/93142/1/EML_18_eng.pdf?ua=1/), datum pristupa: 28.10.2016.
4. Krafs K, Hempelmann E, Skórska-Stania A. From methylene blue to chloroquine: a brief review of the development of an antimalarial therapy. *Prasitol Res.* 2012; 111:1–6.
5. Bhagavathula AS, Elnour AA, Shebab A. Alternatives to currently used antimalarial drugs: in search of a magic bullet. *Infect Dis Poverty.* 2016; 5:103–114.
6. Meissner PE, Mandi G, Coulibaly B, Witte S, Tapsoba T, Mansmann U, Rengelshausen J, Schiek W, Jahn A, Walter-Sack I, Mikus G, Burhenne J, Riedel K-D, Schirmer RH, Kouyaté B, Müller O. Methylene blue for malaria in Africa: results from a dose-finding study in combination with chloroquine. *Malar J.* 2006; 5:84–88.

7. Zoungrana A, Coulibaly B, Sié A, Walter-Sack I, Mockenhaupt FP, Kouyaté B, Schrimmer RH, Klose C, Mansmann U, Meissner P, Müller O. Safety and efficacy of methylene blue combined with artesunate or amodiaquine for uncomplicated falciparum malaria: a randomized controlled trial from Burkina Faso. *PLoS One*. 2008; 3:e1630.
8. Bountogo M, Zoungrana A, Coulibaly B, Klose C, Mansmann U, Mockenhaupt FP, Burhenne J, Mikus G, Walter-Sack I, Schrimmer RH, Sié A, Meissner P, Müller O. Efficacy of methylene blue monotherapy in semi-immune adults with uncomplicated falciparum malaria: a controlled trial in Burkina Faso. *Trop Med Int Health*. 2010; 15: 713–717.
9. Coulibaly B, Pritsch M, Bountogo M, Meissner PE, Nebié E, Klose C, Kieser M, Berens-Riha N, Wieser A, Sirima SB, Breitzkreutz J, Schrimmer RH, Sié A, Mockenhaupt FP, Drakeley C, Bousema T, Müller O. Efficacy and safety of triple combination therapy with artesunate-amodiaquine-methylene blue for falciparum malaria in children: a randomised controlled trial in Burkina Faso. *J Infect Dis*. 2015; 211:689–697.
10. <https://clinicaltrials.gov/ct2/show/NCT01668433?term=methylene+blue&rank=5/>, datum pristupa: 4.11.2016.
11. <https://clinicaltrials.gov/ct2/show/NCT02851108?term=methylene+blue&rank=14/>, datum pristupa 4.11.2016.
12. <https://clinicaltrials.gov/ct2/show/NCT02696928?term=methylene+blue&rank=32/>, datum pristupa 4.11.2016.
13. <https://clinicaltrials.gov/ct2/show/NCT02831023?term=methylene+blue&rank=34/>, datum pristupa 4.11.2016.
14. Congdon EE, Wu JW, Myeku N, Figueroa YH, Herman M, Marinec PS, Gestwicki JE, Dickey CA, Yu WH, Duff KE. Methylthioninium chloride (methylene blue) induces autophagy and attenuates tauopathy *in vitro* and *in vivo*. *Autophagy*. 2012; 8:609–622.
15. Wischik CM, Harrington CR, Storey JMD. Tau-aggregation inhibitor therapy for Alzheimer's disease. *Biochem Pharmacol*. 2014; 88:529–539.
16. <https://clinicaltrials.gov/ct2/show/NCT02245568?term=trx0237&rank=2/>, datum pristupa: 7.11.2016.
17. Rodriguez P, Zhou W, Barrett DW, Altmeyer W, Gutierrez JE, Li J, Lancaster JL, Gonzales-Lima F, Duong TQ. Multimodal randomized functional MR imaging of the effects of methylene blue in the human brain. *Radiology*. 2016; 281:516–528.
18. Atamna H, Nguyen A, Schultz C, Boyle K, Newberry J, Kato H, Ames BN. Methylene blue delays cellular senescence and enhances key mitochondrial biochemical pathways. *FASEB J*. 2008; 22:703–712.
19. Poteet E, Winters A, Yan LJ, Shufelt K, Green KN, Simpkins JW, Wen Y, Yang SH. Neuroprotective actions of methylene blue and its derivatives. *PLoS One*. 2007; 7: e48279.
20. Wen Y, Li W, Poteet EC, Xie L, Tan C, Yan LJ, Ju X, Liu R, Qian H, Marvin MA, Goldberg MS, She H, Mao Z, Simpkins JW, Yang S-H. Alternative mitochondrial electron transfer as a novel strategy for neuroprotection. *J Biol Chem*. 2011; 286:16504–16515.
21. Sontag EM, Lotz GP, Agrawal N, Tran A, Aron R, Yang G, Necula M, Lau A, Finkbeiner S, Glabe C, Marsh JL, Muchowski PJ, Thompson LM. Methylene blue modulates huntingtin aggregation intermediates and is protective in Huntingtin's disease models. *J Neurosci*. 2012; 32:11109–11119.
22. Clifton J II, Leikin JB. Methylene blue. *Am J Ther*. 2003; 10:289–291.



23. <https://www.drugs.com/cdi/urogenic-blue.html>; datum pristupa: 3.11.2016.
24. Moldenhauer Brooks M. Methylene blue as an antidote for cyanide and carbon monoxide poisoning. *Sci Mon.* 1936; 43:585–586.
25. The Merck Index. 13. ed. Whitehouse Station: Merck & Co., Inc., 2001.
26. Bradberry SM. Occupational methaemoglobinaemia. Mechanisms of production, features, diagnosis and management including the use of methylene blue. *Toxicol Rev.* 2003; 22:13–27.
27. Ginimuge PR, Jyothi SD. Methylene blue: revisited. *J Anaesthesiol Clin Pharmacol.* 2010; 26:517–520.
28. Paciullo CA, McMahon Horner D, Hatton KW, Flynn JD. Methylene blue for the treatment of septic shock. *Pharmacotherapy.* 2010; 30:702–715.
29. Ribeiro NAM, Stolf NAG, da Silva Junior AF, Viana VJdC, de Carvalho EN, Athanázio R, Reis MG, de Oliveira SA. Methylene blue use in coronary artery bypass surgery: a prospective randomized study of the hemodynamic and inflammatory responses. *Rev Bras Cir Cardiovasc.* 2004; 19:17–23.
30. Ng BK, Cameron AJ. The role of methylene blue in serotonin syndrome: a systematic review. *Psychosomatics.* 2010; 51:194–200.
31. Aeschlimann T, Cerny T, Küpfer A. Inhibition of (mono)amine oxidase activity and prevention of ifosfamide encephalopathy by methylene blue. *Drug Metab Dispos.* 1996; 24:1336–1339.
32. Dufour C, Grill J, Sabouraud P, Behar C, Munzer M, Motte J, Oberlin O, Paci A, Hartmann O. Ifosfamide induced encephalopathy: 15 observations. *Arc Pediatr.* 2006; 13:140–145.
33. Wagner SJ. Virus inactivation in blood components by photoactive phenothiazine dyes. *Transfus Med Rev.* 2002; 16:61–66.
34. Salah M, Samy N, Fadel M. Methylene blue mediated photodynamic therapy for resistant plaque psoriasis. *J Drugs Dermatol.* 2009; 8:42–49.
35. Tardivo JP, Del Giglio A, Paschoal LH, Baptista MS. New photodynamic therapy protocol to treat AIDS-related Kaposi's sarcoma. *Photomed Laser Surg.* 2006; 24:528–531.
36. Zolfaghari PS, Packer S, Singer M, Nair SP, Bennett J, Street C, Wilson M. *In vivo* killing of *Staphylococcus aureus* using a light-activated antimicrobial agent. *BMC Microbiol.* 2009; 9:27–34.
37. Papin JF, Floyd RA, Dittmer DP. Methylene blue photoinactivation abolishes West Nile virus infectivity *in vivo*. *Antiviral Res.* 2005; 68:84–87.
38. Floyd RA, Schneider, Jr. JE, Dittmer DP. Methylene blue photoinactivation of RNA viruses. *Antiviral Res.* 2004; 61:141–151.
39. Wagner SJ, Skripchenko A, Pugh JC, Suchmann DB, Ijaz MK. Duck hepatitis B photoinactivation by dimethylmethylene blue in RBC suspensions. *Transfusion.* 2001; 41:1154–1158.
40. Müller-Breitkreutz K, Mohr H. Hepatitis C and human immunodeficiency virus RNA degradation by methylene blue/light treatment of human plasma. *J Med Virol.* 1998; 56:239–245.
41. Davies J, Burke D, Olliver JR, Hardie LJ, Wild CP, Routledge MN. Methylene blue but not indigo carmine causes DNA damage to colonocytes *in vitro* and *in vivo* at concentrations used in clinical chromoendoscopy. *Gut.* 2007; 56:155–156.

42. Sturmeijer RG, Wild CP, Hardie LJ. Removal or red light minimizes methylene blue-stimulated DNA damage in oesophageal cells: implications for chromoendoscopy. *Mutagenesis*. 2009; 24:253–258.
43. Mokhlesi B, Leikin JB, Murray P, Corbridge TC. Adult toxicology in critical care: Part II: specific poisonings. *Chest*. 2003; 123:897–922.
44. Harvey JW, Keitt AS. Studies of the efficacy and potential hazards of methylene blue therapy in aniline-induced methaemoglobinaemia. *Br J Haematol*. 1983; 54:29–41.
45. Levin RL, Degrange MA, Bruno GF, Del Mazo CD, Taborda DJ, Griotti JJ, Bouillon FJ. Methylene blue reduces mortality and morbidity in vasoplegic patients after cardiac surgery. *Ann Thorac Surg*. 2004; 77:496–499.
46. Stawicki SP, Sims C, Sarani B, Grossman MD, Gracias VH. Methylene blue and vasoplegia: who, when, and how?. *Mini Rev Med Chem*. 2008; 8:472–490.
47. Xiang J, Xia X, Jiang Y, Leung AW, Wang X, Xu J, Wang P, Yu H, Bai D, Xu C. Apoptosis of ovarian cancer cells induced by methylene blue-mediated sonodynamic action. *Ultrasonics*. 2011; 51:390–395.
48. Wondrak GT. NQO1-activated phenothiazinium redox cyclers for the targeted bioreductive induction of cancer cell apoptosis. *Free Radic Biol Med*. 2007; 43:178–190.
49. Folkes LK, Wardman P. Enhancing the efficacy of photodynamic cancer therapy by radicals from plant auxin (indole-3-acetic acid). *Cancer Res*. 2003; 63:776–779.
50. Kristiansen JE. Dyes, antipsychotic drugs, and antimicrobial activity. Fragments of a development, with special reference to the influence of Paul Ehrlich. *Dan Med Bull*. 1989; 36:178–185.
51. [https://www.genelink.com/Literature/ps/Methylene Blue Oligo staining V2.1.pdf/](https://www.genelink.com/Literature/ps/Methylene%20Blue%20Oligo%20staining%20V2.1.pdf/), datum pristupa: 9.11.2016.
52. Kucsera J, Yarita K, Takeo K. Simple detection method for distinguishing dead and living yeast colonies. *J Microbiol Methods*. 2000; 41:19–21.
53. Christodoulou D, Kandel G, Tsiianos EV, Marcon N. Endoscopic resection of colonic polyps – A review. *Ann Gastroenterol*. 2007; 20:180–194.
54. Simmons RM, Smith SM, Osborne MP. Methylene blue dye as an alternative to isosulfan blue dye for sentinel lymph node localization. *Breast J*. 2001; 7:181–183.
55. Beretvas RI, Ponsky J. Endoscopic marking: an adjunct to laparoscopic gastrointestinal surgery. *Surg Endosc*. 2001; 15:1202–1203.
56. Davies JP, Harris WH. The effect of the addition of methylene blue on the fatigue strength of Simplex P bone-cement. *J Appl Biomater*. 1992; 3:81–85.
57. Oz M, Lorke DE, Hasan M, Petroianu GA. Cellular and molecular actions of methylene blue in the nervous system. *Med Res Rev*. 2011; 31:93–117.
58. [http://www.primehealthchannel.com/sentinel-lymph-node.html#Sentinel Lymph Node Mapping/](http://www.primehealthchannel.com/sentinel-lymph-node.html#Sentinel_Lymph_Node_Mapping/), datum pristupa: 16.1.2017.
59. Skoog DA, West DM, Holler FJ. *Osnove analitičke kemije*. VI. izdanje. Zagreb: Školska knjiga, 1999.
60. Budd MS, Bewick HA. Photometric determination of sulfide and reducible sulfur in alkalies. *Anal Chem*. 1952; 24:1536–1540.
61. [http://www.merckmillipore.com/INTL/en/product/Sulfide-Test,MDA\\_CHEM-114779#documentation/](http://www.merckmillipore.com/INTL/en/product/Sulfide-Test,MDA_CHEM-114779#documentation/), datum pristupa: 14.11.2016.

62. [http://www.kwiaht.org/images/bioremediation/toxicology/MBAS\\_method\\_081408.pdf](http://www.kwiaht.org/images/bioremediation/toxicology/MBAS_method_081408.pdf), datum pristupa: 14.11.2016.
63. <http://www.fann.com/public1/pubsdata/Manuals/Methylene%20Blue%20Kit.pdf>, datum pristupa: 3.11.2016.
64. [http://stoltz.caltech.edu/seminars/2008\\_Winston.pdf](http://stoltz.caltech.edu/seminars/2008_Winston.pdf), datum pristupa: 14.11.2016.
65. Boyd JF, Simmonds RC. Continuous laboratory production of fertile *Fundulus heteroliticus* (Walbaum) eggs lacking chorionic fibrils. J Fish Biol. 1974; 6:389–394.

Primljeno 16. studenoga 2016.